



## DEMENTIA CARE SERIES

Michigan Department of Health and Human Services

CARING SHEETS: THOUGHTS & SUGGESTIONS FOR CARING

# Caring Sheet #11: Alzheimer's Disease:

A Summary of Information and Intervention Suggestions with an Emphasis on Cognition

By Shelly E. Weaverdyck, PhD

## Introduction

This caring sheet focuses on Alzheimer's Disease, with an emphasis on cognition.

Caring sheets #12 and #13 summarize information and intervention suggestions regarding Dementia with Lewy Bodies (caring sheet #12) and Frontotemporal Dementia (caring sheet #13). All three outline the brain changes in each type of dementia, the impact these changes have on cognition and behavior, and implications for effective intervention.

The three caring sheets (#11, #12, and #13) are written as companion pieces in outline form with virtually each line of one caring sheet corresponding with each line of the other two. The three can be laid out side by side, and compared almost line by line.

Other caring sheets describe the cognitive and behavioral changes and specific interventions for these dementias in more detail. Caring Sheet #2 in particular describes the characteristics of dementia, and the relationship between the brain changes and changes in cognition.

Dementia is a decline in a person's cognition. This decline occurs because of changes in the brain.

If the cognitive decline is caused by treatable disorders such as a urinary tract infection, vitamin deficiency, reactions to medications, or depression, it is likely **temporary** and **treatable** (e.g., delirium).

In other cases the brain changes and resulting cognitive decline are **irreversible** and **progressive** (i.e., increasingly severe). They are caused by disorders such as Alzheimer's Disease, vascular disorders (e.g., ministrokes), Creutzfeldt-Jakob Disease, Dementia with Lewy Bodies, or Frontotemporal disorders. There are over 80 different disorders that cause this type of progressive dementia. Alzheimer's Disease is the most common.



## **ALZHEIMER'S DISEASE (AD)**

### **CHARACTERISTICS**

Brain disorder

Most obvious symptoms: memory loss and cognitive impairment

Progression: increasing severity of symptoms over time (a progressive dementia)

Onset: insidious; age 40-90, usually over age 65 (median age 73.5)

Duration: average 8 years from onset to death (may vary with time of diagnosis)

Cause unknown

Cure: no cure at this time, but there is treatment to reduce symptoms

Diagnosis verified at autopsy

Is the most common cause of dementia (60% of all dementia cases)

Affects 10% of all people over age 65

Risk factors: age, APOe4 gene, Down's Syndrome, family history

Hereditary in 10% of cases

Course: gradual, steady decline, decline 2-4 points per year on Folstein Mini-Mental State Exam, no spontaneous improvement

Alois Alzheimer first described neuropathology in a 51-year-old woman in 1907

### **NEUROPATHOLOGY**

Neuritic plaques outside of cells in brain

Neurofibrillary tangles inside of cells in brain

Atrophy (i.e., loss) of brain tissue; cell death

Acetylcholine reduction

### **LOCATION OF CORTICAL BRAIN CHANGES**

Cortical refers to the cortex (i.e., the outer layer) of the brain

Changes (pathological abnormalities) occur in the cortex and in internal (subcortical) structures of the brain

Changes (pathological abnormalities) occur on both sides of the brain

Order of cortical brain structures affected, creating stages:

Hippocampus (subcortical)

Parietal lobes

Temporal lobes (posterior then anterior)

Frontal lobe



## **COGNITIVE CHANGES**

Memory impairment first obvious symptom

Visuospatial perception and skill impairment:

    Difficulty recognizing distance between objects and from self

Difficulty arranging objects in space

Language impairment:

    Difficulty understanding and producing speech

    Difficulty reading and writing

    Difficulty understanding what is being read, when can read

Insight impaired

Judgment impaired

Disorientation

Concentration impaired

Abstract processing impaired

Attention impaired

Sense of time impaired

Ability to analyze, plan, organize impaired

No sensory loss (though may occur with normal aging)

No focal deficits

No disturbance of consciousness

Person becomes increasingly dependent on environment throughout course

## **EMOTIONAL CHANGES**

Depression throughout course, especially in early stages

Emotional lability (switch quickly from one emotion to another)

Less expression of emotional intensity and switching in later stages

## **BEHAVIORAL CHANGES**

Get lost occasionally in early stages; increasingly often throughout course

Loses objects occasionally in early stages; increasingly often throughout course

Distressing behaviors usually triggered by anxiety, confusion and  
    misinterpretation of environment in middle stages

Distressing behaviors usually triggered by physical pain and discomfort in later stages

Gait and physical movements preserved until later stages

Incontinence only in later stages

Hallucinations when they occur, occur in later stages



### **INTERVENTIONS: Non-medicinal**

- Assess individual regularly over time as dementia progresses
- Modify expectations and intervention as change occurs
- Express warmth and affection for person, verbally and nonverbally
- Respect person as adult; avoid treating person as a child
- Stay calm and help the person feel relaxed
- Compensate for the person's cognitive impairment to help person feel good and comfortable and to prevent fatigue and embarrassment
- Assess and modify environment, caregiver interactions and daily routines over time throughout course
- Maintain appropriate stimulation that engages person
- Avoid meaningless or confusing stimulation
- Use touch as appropriate when communicating
- Simplify daily routines and tasks
- Provide information through cues in environment and when communicating
- Ensure cues are understandable to person
- Avoid arguing with the person
- Address person's feelings and then distract when person engaged in distressing behavior
- Explain to person what caregiver is doing and intends to do
- Use concrete requests and concrete cues
- Compensate for sensory changes that occur with normal aging

### **MEDICAL TREATMENTS**

- Cure unknown
- Reduce loss of acetylcholine (inhibit acetylcholinesterase)
  - Cognex, Aricept, Exelon, Reminyl
  - Most effective in mild-moderate stages
- Memantine (regulates activity of glutamate)
  - Can be effective in severe stages (for persons with MMSE score of 3-14)
- Cholinergic medications for behavioral symptoms
- Antioxidants
- Anti-inflammatory agents
- Neurotrophic compounds
- Anti-amyloid deposition
- Vitamin E in high doses (only under doctor's care) (effectiveness is controversial)



To extent AD associated with cardiovascular risk factors, can prevent by reducing cardiovascular risk factors: Exercise, Diet, Cholesterol-lowering medications (statins) such as Lipitor

## **COMMENTS**

In 1984 criteria for clinical diagnosis created

Red flags that suggest a disorder is probably not AD:

Onset before age 60 years

Sudden onset

Rapid progression

Symptoms that do not occur in the order of typical AD stages

Behavior changes or hallucinations occur much earlier than  
memory impairment

Incontinence occurs before later stages

Seizures occur before later stages

(Exception: Downs may cause earlier seizures)

Abnormal neurological symptoms (impaired gait, falls, weakness) occur before  
later stages

## **RESOURCES**

<http://www.ninds.nih.gov/disorders/alzheimersdisease/alzheimersdisease.htm>

(National Institute of Neurological Disorders and Stroke NINDS)

<http://www.alzheimers.org/> (Alzheimer's Disease Education and Referral Center  
ADEAR)

<http://www.alz.org/> (Alzheimer's Association)

<http://www.med.umich.edu/madrc/> (Michigan Alzheimer's Disease Research Center  
MADRC)



This caring sheet focuses on Alzheimer's Disease (AD). More details about the brain changes and resulting cognitive changes in AD are in caring sheet #2.

Alzheimer's Disease is by far the most common cause of irreversible brain changes & dementia in persons over the age of 65 years.

Though these changes in behavior and cognition result from brain damage, they are often mistakenly viewed as intentional or manipulative.

In AD the four hallmark pathological changes to the brain are:

- **Atrophy:** the reduction in size of a structure. Atrophy due to death of nerve cells in AD causes much of the confusion and cognitive impairment. Atrophy occurs with normal aging, but is especially pronounced and is pathological in AD. The atrophy is visible on a CAT scan and at autopsy.
- **Neuritic plaques:** little patches or collections of debris in the brain. They are located outside of nerve cells. A protein called amyloid is at the core of the plaques. The number of neuritic plaques correlates with a person's performance on cognitive tests (such as Intelligence or IQ tests).
- **Neurofibrillary tangles:** inside nerve cells, particularly in the axons of nerve cells. Axons are the protrusions from the nerve cell body, which carry information from one nerve cell to the next. Tiny neurofibrils (filaments or tubules) transport cell nutrients within the nerve cell. The neurofibrils become tangled in a very characteristic way (double helical) and therefore disrupt the cell maintenance processes, probably contributing to the cell's death. The number of neurofibrillary tangles also correlates with a person's performance on various cognitive tests.
- **Neurochemical changes:** a reduction in some of the neurotransmitters. Neurotransmitters (the chemical messengers) are neurochemicals which are transferred from one nerve cell to another as a method of communication with that nerve cell. This intercellular communication is essential to the brain's maintenance and functioning. There are many different kinds of neurotransmitters. One, which is particularly reduced in amount in Alzheimer's Disease, is called acetylcholine.



## Vascular Dementia

Vascular dementia is another common disorder. In Vascular dementia the brain has many tiny strokes on the cortex or surface of the brain (as well as other areas). Spots of softened dead tissue (lesions) occur throughout the cortex. These lesions can sometimes be seen on a CAT scan. A CAT scan can rule out the possibility of a major stroke as a source of the cognitive impairment. The cognitive changes are more varied than they are in AD.

## Other Dementias

See caring sheet #12 for more information about Dementia with Lewy Bodies (DLB) and caring sheet #13 for more information about Frontotemporal Dementia (FTD). DLB is another more common disorder. FTD is less common but can be a particularly challenging disorder.

© Copyright 2005 by S. Weaverdyck (Page 7 and Header Revised 2018).

Citation: Weaverdyck, S. (2005). #11 - Alzheimer's Disease: A Summary of Information and Intervention Suggestions with an Emphasis on Cognition. *Dementia Care Series Caring Sheets: Thoughts & Suggestions for Caring*. Lansing, Michigan: Michigan Department of Health and Human Services.

Edited and produced by Eastern Michigan University (EMU) Alzheimer's Education and Research Program for the Michigan Department of Health and Human Services (MDHHS), with gratitude to the Huron Woods Residential Dementia Unit at St. Joseph Mercy Hospital, Ann Arbor, Michigan.

The author, Shelly Weaverdyck was Director of the EMU Alzheimer's Education and Research Program.

Editor: Shelly Weaverdyck, PhD, Former Director EMU Alzheimer's Education and Research Program; Email: sweaverd@umich.edu

All Caring Sheets are available online at the following websites:

**[http://www.michigan.gov/mdhhs/0,5885,7-339-71550\\_2941\\_4868\\_38495\\_38498---,00.html](http://www.michigan.gov/mdhhs/0,5885,7-339-71550_2941_4868_38495_38498---,00.html)** (Michigan Department of Health and Human Services MDHHS), at **<http://www.lcc.edu/mhap>** (Mental Health and Aging Project (MHAP) of Michigan at Lansing Community College in Lansing, Michigan), and at **<https://www.improvingmipractices.org/populations/older-adults>** (Improving MI Practices website by MDHHS)

The Caring Sheets were originally produced as part of the in-kind funding for the Michigan Alzheimer's Demonstration Project. Funded by the Public Health Service, Health Resources and Services Administration (1992-1998) and the Administration on Aging (1998-2001) 55% federal funding and 45% in-kind match. Federal Community Mental Health Block Grant funding supported revisions to Caring Sheets (2002-2018).